



CIRCUMCISION IN LONDON

Forceps Guided (traditional) method - Children




Your son can walk normally.
No running, jumping or physical activity for at least 2 weeks



Apply Vaseline as we show you for 2 weeks



Shower your son daily only after 3 days. Dry the penis after the shower.



Sometimes we place stitches. They dissolve in 1-4 weeks. Don't worry if they open early, the wound will close by itself.



Swelling & redness is normal.



Yellowish/ black crust is normal & will exfoliate naturally. This is NOT an infection






Patient older than 2 months can be given Calpol (paracetamol). Always read the leaflet



A few drops of blood is normal. If there is continuous bleeding, please contact us.

Pain – your son maybe unsettled after the circumcision which is normal. This will vary from patient to patient and can last for several hours to several days. If your son is older than 2 months, you can give Calpol only (Always read the leaflet).

If the head of the penis is covered: you must press down the pubic area and pull back the shaft skin to fully expose the head of the penis daily to clean and dry. You may also need to pull out the head of the penis when doing this. This appearance is due to swelling or buried penis.



Scan on your phone camera to see an explanation video

Dr S Al-Ali & Dr H Al-Ali
Beverley Clinic
78 Beverley Drive
Edgware
HA8 5NE

020 8951 3794
0788 6779958

info@circumcisioninlondon.co.uk
www.CircumcisionInLondon.co.uk

Continuous bleeding emergencies only:

Please apply pressure as shown in the video in the above QR code. If the bleeding does not stop, please call both clinic numbers to the left and listen to the whole answer machine message to obtain the emergency number. In the rare event of not reaching us, please call 999.